

## WEST BENGAL STATE UNIVERSITY

B.Sc. Honours 2nd Semester Examination, 2019

# CEMACOR03T-CHEMISTRY (CC3)

Time Allotted: 2 Hours

1



Full Marks: 40

The figures in the margin indicate full marks. Candidates should answer in their own words and adhere to the word limit as practicable. All symbols are of usual significance.

### Answer any four questions taking one from each unit

#### <u>Unit-I</u>

1.	(a)	Energy of an excited electron of $He^+$ ion is -6.04 eV. Calculate the angular momentum of that excited electron from Bohr's theory.	4
		[Given: Energy of 1st Bohr Orbit of Hydrogen $= -21.8 \times 10^{-19}$ J.	
		$1 \text{eV} = 1.6 \times 10^{-19} \text{ J}, \ h = 6.6203 \times 10^{-34} \text{ Js}$	
	(b)	Calculate the wave length of spectral line when the electron jumps from the third to the second Bohr orbit in a hydrogen atom.	4
		[Given: Rydberg constant = $109677 \text{ cm}^{-1}$ ]	
	(c)	Using $p^2$ electronic configuration determine number of microstates. Write symbol of Ground state.	3
2.	(a)	Frame an equation for the energy of an electron revolving in a circular path around hydrogen atom.	4
	(b)	Calculate the Principal quantum number where the electron is revolving in a H-atom, when its kinetic energy is $217.945 \times 10^{-20}$ J.	2
		[Given: $E_1$ for $H = -871.78 \times 10^{-20} \text{ J}$ ]	
	(c)	Which set of orbitals is defined by the quantum numbers $n = 3$ and $l = 2$ ? How many orbitals will be there?	2
	(d)	Find out the spectroscopic ground state term symbols for $Cu^{2+}$ and $Cr^{2+}$ ions.	3

#### <u>Unit-II</u>

3.	(a)	The ionisation potential of neon is 21.56 eV while that of the next element, sodium	2
		is only 5.14 eV. — Explain.	
	(b)	The atomic radii of Zr and Hf are almost same — Explain.	3
	(c)	Which of the following reactions should proceed spontaneously in the gas phase?	3
		Explain with reason.	

(i) 
$$Xe + He^+ \longrightarrow Xe^+ + He$$

(ii) 
$$Si + Cl^+ \longrightarrow Si^+ + Cl$$

(iii) 
$$F^- + I \longrightarrow F^- + I^-$$

#### CBCS/B.Sc./Hons./2nd Sem./Chemistry/CEMACOR03T/2019

C	SC 5/	<b>B.Sc./Hous./Zhu Sem./Chemistry/CE/WACOR051/2019</b>			
4.	. (a)	Explain when third ionization energy (IE <sub>3</sub> ) of Mg (7732.8 kJ/mole) is about ten times higher than that of first ionization energy (IE <sub>1</sub> , 737.7 kJ/mole).	3		
	(b)	The second electron attachment enthalpy of oxygen is positive — yet it forms most compounds in $-2$ oxidation state — Justify.	3		
	(c)	PbCl <sub>4</sub> is very unstable and oxidising — Why?	2		
<u>Unit-111</u>					
5.	(a)	What is meant by buffer solution? Give an example of an acidic and a basic buffer solution.	2+2=4		
	(b)	State Bronsted-Lowry concept of acids and bases. Write down the conjugate acid/base of the following:	2+2=4		
		$H_2CO_3$ , $SO_4^{2-}$ , $NH_4^+$ and $CH_3COOH$			
	(c)	What are super acids? Give one example.	2		
6.	(a)	Name the indicator you would select for the titration of aqueous solutions of $Na_2CO_3$ and $H_2SO_4$ . Give reason in brief.	3		
	(b)	Calculate the pH of an aqueous solution of ammonium acetate at 25°C. [Given: Dissociation constant of acetic acid, $K_a = 1.78 \times 10^{-5}$ and that of	4		
		ammonium hydroxide is $1.8 \times 10^{-5}$ .]			
	(c)	Explain solvent levelling and differentiating effects.	3		
		<u>Unit-IV</u>			
7.	(a)	Balance the following equations by ion-electron method: (i) $NaBiO_3 + Mn(NO_3)_2 + HNO_3 \longrightarrow Bi(NO_3)_2 + NaNO_3 + H_2O + NaMnO_4$ (ii) $K_2Cr_2O_7 + KI + HCl \longrightarrow CrCl_3 + KCl + I_2 + H_2O$	2+2=4		
	(b)	$E^{0}$ of Ce <sup>4+</sup> /Ce <sup>3+</sup> couple is 1.44 V. Find <i>E</i> at equivalence point in the titration of 0.1 N Fe <sup>2+</sup> by 0.1 N Ce <sup>4+</sup> in 1M acid medium ( $E^{0}$ for Fe <sup>3+</sup> /Fe <sup>2+</sup> couple = 0.77 V).	3		
	(c)	Calculate the solubility of $CaF_2$ in solutions of	4		
		(i) $10^{-3}$ M Ca(NO <sub>3</sub> ) <sub>2</sub> (ii) $10^{-1}$ M NaF [ $K_s = 1.7 \times 10^{-10}$ ].			
8.	(a)	Establish the Nernst equation for the couple $VO_3^-/VO^{2+}$ . If the $E^0$ value for the couple in 1M acid medium at 25°C be 0.92 volt, calculate its formal potential at pH 7.	1+2=3		
	(b)	From the reduction potentials of the following sequences of reactions predict whether $MnO_4^-$ or $MnO_4^{2-}$ is the better oxidant if the reduction product is $Mn^{2+}$ .	2		
		$MnO_{4}^{-} \xrightarrow{0.56 \text{ Volt}} MnO_{4}^{2-} \xrightarrow{2.26 \text{ Volt}} MnO_{2} \xrightarrow{0.95 \text{ Volt}} Mn^{3+} \xrightarrow{1.51 \text{ Volt}} Mn^{2+}$			
	(c)	$Cu^{2+}$ is unstable in aqueous medium and undergoes disproportionation reaction. — Explain. [Given: $E_{Cu^{2+}/Cu^{+}}^{0} = 0.15$ V, $E_{Cu^{2+}/Cu^{0}}^{0} = 0.34$ V]	3		
	(d)	Discuss the solubility product principle. Explain how this principle is utilised in the precipitation of Gr-IIIB ( $Co^{++}$ , $Ni^{++}$ , $Mn^{++}$ and $Zn^{++}$ ) cations as sulphides in qualitative analysis.	3		

2009

X